

CARE FOLLOWING DENTAL PROCEDURES

Proper care of the mouth following most dental procedures can reduce complications and speed the healing of the surgical area.

- Protection of blood clot** . . . maintain gentle pressure by biting on the gauze sponge that has been placed over the surgical area, or by biting on a tea bag which has been gently moistened and wrapped in a piece of gauze. Keep steady firm pressure for 45 minutes. Repeat as often as needed.
- Do not rinse** . . . or use a mouth wash for at least 24 hours. After 24 hours rinse with warm salt water (½ teaspoon table salt in 8 oz. of warm water) every 1–2 hours is recommended. (The use of commercial mouth washes during the healing period is not encouraged.)
- Discomfort** . . . following dental surgery it is normal to experience some discomfort. If medication has been given or prescribed, take as instructed.
- The toothbrush** . . . may be carefully used in the area of the mouth not involved by the surgical procedures. A clean mouth heals faster.
- Eating** . . . adequate food and fluid intake following surgery and/or general extractions is most important. If you find that eating your regular diet is too difficult you may supplement your diet with liquids such as Carnation Instant Breakfast (CIB/any flavor). The high quality, nutritional value of CIB is important for prompt healing following such procedures. Should you not be able to chew solid foods of any nature, for several days or longer, follow the nutritionally balanced liquid diet on the reverse side.
- Avoid** . . . all excessive activity, don't pick at the surgical area, don't consume liquids through a straw, avoid alcoholic beverages and refrain from smoking until healing is well established.
- Sutures** . . . if they were used, do not fail to return for their removal on the appointment date given.
- Control of swelling** . . . gently apply ice packs to area for periods of 20 minutes on, 10 minutes off. This procedure should continue for the first 24 hours only.
- Allergic reactions** . . . for generalized rash, itching, etc., call dentist immediately.
- Do not hesitate** . . . to call if any questions arise.

OTHER INSTRUCTIONS

MICHAEL I. LEVY, D.D.S.
general dentistry



Phone (602) 274-9242
e-mail fenixflea@aol.com

59 E. Lexington
Phoenix, AZ 85012
1 Blk. South of Osborn
at 2nd Street

“ENSURE”

SEMI-LIQUID DENTAL DIET

If a semi-liquid diet should be necessary for several days, the following diet has been designed to assist you in selecting meals which supply ample amounts of essential nutrients. The sample menu is approximately 2,000 calories and supplies adequate quantities of essential nutrients recommended by the Nutrition Board of the National Research Council, 1980.

Basic Meal Plan	Sample Menu
BREAKFAST Fruit Juice CIB* mixed with milk beverage Cooked Cereal Sweets (optional) Beverage	6 oz. orange juice 1 serving of CIB* made with nonfat milk 1 cup Instant Farina 1 tsp. honey Coffee or tea
LUNCH Soup Starch Smooth Entree Fruit CIB* mixed with milk beverage	6 oz. vegetarian vegetable soup 1 packet (4 crackers) Saltines (dipped in the soup to soften) 1 cup lowfat cottage cheese served with ½ medium banana and ½ canned peach 1 serving of CIB* made with nonfat milk
DINNER Starch Smooth Entree Vegetable Dessert Milk Beverage	1 cup cooked spaghetti noodles 4 oz. lean ground beef in ½ cup tomato sauce ½ cup mashed winter squash ½ cup pudding made with milk 8 oz. nonfat milk
BEDTIME SNACK CIB* mixed with milk beverage	1 serving of CIB* made with nonfat milk

Entrees may consist of any flaked, shredded, ground, or pureed meats, fish or fowl mixed with smooth sauces or gravies, or cheese or egg dishes.

Fruits and vegetables should be cooked, canned, pureed, or souffled. Desserts include custards, puddings, junkets, and ice cream.

*CIB/ Carnation Instant Breakfast—any flavor.